

PEG-ELECTROLYTE SOLUTION  
BOWEL PREPARATION FOR **COLONOSCOPY AT OR BEFORE 10:00AM**

These instructions have been prepared to help you understand the procedure for cleansing the colon (large intestine). A thorough examination depends on your colon being carefully cleansed and emptied. Stool remaining in the colon can obscure important details, resulting in the exam having to be repeated at another time.

Your Colonoscopy Appointment is on (date) \_\_\_\_\_ at (time) \_\_\_\_\_ at (location) \_\_\_\_\_

**READ THIS ENTIRE INSTRUCTION SHEET**

**On The Day Before Your Colonoscopy:**

1. **Be sure you have arranged a driver who will stay with you and drive you home. Please inform your driver this will take about 2 hours.**
2. **NO FOOD** on the day before your procedure. Follow the clear liquid dietary instructions. (See clear liquid diet list at right.)
3. If you are experiencing **constipation**, take two (2) Dulcolax tablets at **12:00 PM (Noon)**.
4. At **6:00 PM**, begin drinking the Preparation Solution. Drink one glassful (8 oz.) every 5-10 minutes, or as fast as tolerated, **until the gallon is gone**. The prep often works within 30-60 minutes. Remain close to the toilet, as multiple bowel movements will occur. To help with the taste, you may find it helpful to add to **each glass** some fresh lemon juice, Crystal Light™ powder, sugar-free Kool-aid™ powder, or sugar substitute. **Do not add any product containing sugar as this may alter the quality of the prep.**
5. After completing the preparation solution, continue the **CLEAR LIQUID** diet.
6. You should be passing clear or light yellow fluid.

**\*\*\*If you are unable to complete this preparation call the number listed on the right side of the sheet.\*\*\***

**On The Morning of Your Colonoscopy:**

1. **NO FOOD**. You may continue with clear liquids until **3 hours** prior to your colonoscopy appointment. After that nothing by mouth until after your exam.
2. If you regularly take medication in the morning, take it at least **3 hours** before your colonoscopy appointment, with a small amount of water.

**Medications You Need to Cleanse the Bowel:**

- “Preparation Solution” (Golytely, Nulytely, Colyte or Trilyte) 1 gallon.
- **Dulcolax 5 mg** (bisacodyl) 2 tablets. This is an over the counter medication, no prescription required. (Use **only** if you are experiencing constipation.)
- ✓ **Your “Preparation Solution” prescription has been sent electronically to your pharmacy unless otherwise mentioned.**
- ✓ **ONLY FOLLOW THE DIGESTIVE HEALTH SPECIALISTS INSTRUCTION SHEET (DO NOT FOLLOW THE DIETARY INSTRUCTIONS GIVEN BY THE PHARMACY)**

**Clear Liquid Diet List:** *(Any liquid that is clear enough to read print through it.)*

- Clear fruit juices without pulp (apple, white grape)
- Water, tea or black coffee
- Low sodium **CLEAR** broths
- Jell-O™ – lemon, lime or orange, without topping or fruits
- Popsicles™
- Kool-aid™, Tang™, Crystal Light™, Gatorade™, soda pop
- **NO FOOD, NO FRUIT PIECES, NO PULP, NO RED FLAVORS OR DYES**
- **NO ALCOHOL**

**Some Suggestions While Prepping At Home:**

- You may experience chills. Dress warmly. (socks, robe, blanket)
- Walking around the house may help get the prep working.
- Drinking from a straw may be helpful.
- If you experience **nausea or vomiting**, stop drinking for 1 hour and then start again at a slower rate. Drink each glass in 10-20 minutes, resting 10 minutes between each glass.
- Be patient, the prep solution rarely fails. Most patients have a bowel movement within 30-60 minutes, but it may take up to **two hours**.

**24-Hour Phone Line:**

**Tacoma: 253-272-5127 • Federal Way: 253-838-9839**  
**Puyallup, Downtown: 253-841-3933 • Gig Harbor: 253-858-0112**  
**Puyallup, Sunrise at South Hill: 253-770-3700**

Please visit us at: [www.dhsgi.com](http://www.dhsgi.com)